OPEN SOURCE INVESTIGATION

IDENTIFICATION OF ECOLOGICALLY FRAGILE AREAS PRIORITIZED FOR FOREST CONSERVATION AND RESTORATION IN MAMBASA AND MONGBWALU SECTOR, ITURI PROVINCE, DEMOCRATIC REPUBLIC OF THE CONGO









watetezi.org

Environmental Defenders, March 2021

Jonam Hill-Dei PO-BOX, 9520 Kampala, Uganda



with the technical contribution of COMUNI-DADOS



Open Source Investigation

FORÊT DE SANG

Identification of ecologically fragile areas prioritized for forest conservation and restoration in Mambasa and Mongbwalu sector, Ituri province, Democratic Republic of the Congo

WEBGIS: http://watetezi.org/data/DRC/analysis.html



The restoration of degraded ecosystems is an essential activity to achieve environmental and ecological justice. The general approach is to implement actions aimed at conserving intact ecosystems as they provide ecosystem services that are not fully recoverable upon destruction and to restore degraded ecological and ecosystem systems because degradation is taking place faster than recovery. An ecosystem is defined as restored when following a self organized ecological trajectory (change over time in ecological characteristics) in line with the desired or reference ecosystem. Implementing conservation and restoration projects

by evaluating them only from a biological perspective is not enough, as they must ensure the inclusion of local communities through a rights-based approach. Environmental and ecological justice focus on the intersections between the systemic exploitation of humans and the natural world; this includes inequities in the access and use of natural resources and in the distribution of environmental harms (by race, class, gender, among many). Furthermore, land rights for women in Africa go beyond property rights and touch on often sensitive issues around different tenure systems (statutory, customary and religious), landbased wealth, power and social relations that give or take away their right to access and control resources. These kinds of projects fit neatly within the perspective of ecological politics through whose lens we question the viability of extractive and food production models, providing alternatives to the ecological, economic, and democratic crises plaguing the Albertine rift region. Although this perspective is being implemented in a growing number of projects spread across all areas of the planet that face critical conservation issues, some local factors are still crucial in determining the success of these interventions. For this reason, assimilating the local perspective requires a dedicated suitability analysis for the area in which to prioritize and implement conservation and restoration projects (through passive and active methods).

This study focuses on two areas located in the Ituri province, in the portion of the Albertine rift located within the territory of the Democratic Republic of Congo, and flagged as territories strongly impacted by ecological injustice. 3

STUDY AREA

MAMBASA

The landscape of the Mambasa sector is facing a major transformation of its land cover since two decades. It is undergoing a gradual decline in dense and secondary forests, triggered by (legal, illegal and unregulated) resource exploitation and their unsustainable use, some detrimental forms of agriculture, rapid urban growth in absence of long-term strategic planning, all linked by instability arising from land conflicts that chronically plague the region.

The sector being considered is at the latitude/ longitude [676226.57, 95245.37 : 763251.47, 174895.64 EPSG:32635 WGS 84 / UTM zone 35N], and is located southeast of the Okapi Integral Forest Reserve. In particular, the territory south of the city of Mambasa and enclosed between the dense forest and the road to the city of Biakato is analyzed. This area is known for the presence of a high number of artisanal and small-scale mines, along with intense illegal logging operations that have resulted in isolated penetrations into the forest that are causing the fragmentation of the habitat. The expansion of agriculture linked to the increase in population (in particular along the road between Mambasa and Biakanto) and the growth of new urban centralities, are pressing on the ecological system in particular along the road between Mambasa and Nia-Nia that crosses the Okapi forest. Models, project a loss of secondary forest area of up to 13.65% by 2035 in Mambasa sector, but could be even worse considering changes in the composition of biodiversity and the effects of climate change.

Within the resource exploitation scenario in Mambasa, community rights and individual opportunistic interests increasingly clash since the politico-militarycommercial elite seizes the benefits provided by the vacuum of the state and the richness of the rainforest. The composition of the population of Mambasa differs profoundly from other territories in Ituri province. Bantu constitute the majority (70 percent), next to a large group (around 30 percent) of Mbuti or pygmies. Furthermore, there is an influx of Nande from the Beni-Butembo area (North Kivu province). Pygmies are most vulnerable as they have not been able to secure their land rights. But other population groups are also affected as communities and individual customary chiefs are being played against each other.

Given the evidence presented, this sector is targeted for environmental conservation projects involving community-driven right-based processes. Beside that, the implementation of forest restoration projects in buffer zones around vulnerable resources and their sustainable management system to control development proves significant. ITURI PROVINCE FOREST COVER LOSS 2001-2019

Okapis

** ()

, Kibali-Ituri http://watetezi.org/data/DRC/analysis.html Bondon o Manufelia

Open the webmap to navigate and select the information

Kibali-Itari Kibali Tari

Figure 1. Mambasa sector





MONGBWALU

The sector being considered is at the latitude/ longitude [791384.82, 151118.60 : 873908.43, 230264.77 EPSG:32635 WGS 84 / UTM zone 35N], and is located between the city of Mongbwalu and Bunia, the capital city of Ituri province. 6

The small town of Mongbwalu is known for being an industrial-scale gold mining area, where international companies have been present since 1900 (gold was first discovered in the area by Australian prospectors in 1905 and has been mined ever since). Competition over control of these goldfields began during the 1998 Congolese war and is continuing to the present; despite the international community has not officially recognized a link between the exploitation of resources and the armed conflict in Democratic Republic of Congo, a long series of human rights violations have been recorded in the mining sites. In addition, the relationship between national and international companies controlling gold reserves with armed groups through exchanges of logistical and financial support in return for political benefits and security for the workers has been proven (e.g. the AngloGold Ashanti gold producer with the Nationalist and Integrationist Front - FNI armed group). Gold mining operation employs thousand workers and is one of the main providers of employment; people of different ethnicities have come to live in Mongbwalu to work in the gold mines or in related activities.

A century of exploration and exploitation of subsoil resources have seriously compromised the ecological balance of the sector. Therefore, considering the trend of land degradation drivers that have already been critical for years, this sector is targeted for environmental restoration projects, particularly in depleted mining sites and those that have been subject to exploration.

METHODOLOGY

In the study, a GIS based Weighted Overlay Process is applied to identify areas affected by degradation (where to target conservation efforts and activating forest restoration projects); Weighted Overlay is a method of modeling suitability.

Recognition of the suitable areas to the proposal requires many different elements to be evaluated together in the decision making process, and Geographical Information Systems provide convenience for the interpretation of large-scale and complex data.

We call parameters the thematic classes employed to perform a weighted analysis among the elements considered (e.g. biodiversity and natural resources or drivers of conflict). Following that, specific criteria and sub-criteria are hierarchically identified for each of the parameters (e.g. biodiversity intactness index and species range-size are criteria subject to the biodiversity and natural resources parameter) and scored to a common suitability scale range (Fig. 3). The parameters and criteria are identified as a result of the literature researches and studies made specifically for the study area. Weights for determined parameters and scores for criteria are obtained by a participation process involving Civil Society Organizations, community leaders, and with the help of experts. At this stage, complementary methods such as the Analytical Hierarchy Process (AHP) have not been adopted, given the lack of participation in the selection of criteria and parameters of stakeholders who should critically contribute to decision making processes.



WEIGHTED OVERLAY ANALYSIS

Selection of parameters and weights

Six parameters are identified: biodiversity and natural resources; carbon cycle; people; drivers of conflict – mining sector; drivers of conflict – others; alerts. To assign a relative degree of importance (weight) to every parameter based on a fundamental suitability scale, Environmental Defenders collaboratively defined these values through dialogue with the Civil Society Organizations operating in the Ituri province, the systematization of knowledge accumulated on the ground, and the review of the literature (Tab. 1).

Parameter Maps

Parameter: BIODIVERSITY AND NATURAL RESOURCES

Criteria: Species Range-Size Rarity

The Species Range-Size Rarity is calculated as the rarity of species occupancy among areas. In this case, all threatened amphibians, birds and mammals included among IUCN Red List's categories CR (Critically Endangered), EN (Endangered), and VU (Vulnerable) are considered.

Criteria: Species Richness

The Species Richness represents the number of species potentially occurring in a spatial grid. In this case, all threatened amphibians, birds and mammals included among IUCN Red List's categories CR (Critically Endangered), EN (Endangered), and VU (Vulnerable) are considered.

Criteria: Biodiversity Intactness Index

The Biodiversity Intactness Index (BII) estimates how the average abundance of native terrestrial species in a region compares with their abundances before pronounced human impacts. A value around 50% indicates that the species originally present are on average only half as abundant compared with pristine conditions (pre-industrial climate period).

For the purpose of the analysis, is determined a range of BII values within which the impact is defined as critical considering the trend of land use/land cover change and dynamics in the territory over the last ten years.

Criteria: Waterway

Deforestation and associated water pollution threaten the waterways of Ituri Province. Water quality deterioration from heavy metal pollution is a major issue of concern nearby mining sites, particularly given the considerable environmental legacy of intensive industrial-scale mining and with different magnitudes given the artisanal and small-scale mining. Also, the extensive land cover and land use changes encountered in some case studies, have led to excessive rates of soil erosion within many catchments increasing the risk for sediment pollution.

For this reason, just for the scope of this analysis, all areas pertaining to waterways are considered as critical and designated as priorities for protection and conservation (a buffer zone has been defined).

Figure 3 Parameters and Criteria

Criteria: Protected Area

7

Protected areas are established in the state forest domain or in other sites of national, provincial or local interest and include: 1) Natural Integral Reserves; 2) National Parks; 3) Natural monuments; 4) Habitat or species management areas; 5) Biosphere Reserves; 6) Protected land or sea landscapes; 7) Zoological and Botanical Gardens; 8) Hunting Areas and Reserves; 9) any other category that special laws and regulations designate as such for the conservation of species of fauna and flora, soil, water, mountains or other natural habitats. These areas are managed mainly by Law No. 14/003 of 11 February 2014 on nature conservation.

The most representative protected area in the Ituri province is the Okapi Natural Integral Reserve, established in 1992, and listed also as a World Heritage Site (2006). Over the past years, the park has been suffering from increased poaching, deforestation, and illegal mining operations located particularly on the edge of the reserve designated border, but also rebel activity and immigration. For this reason, a 1 km buffer zone surrounding the designated protected area is also considered critical for conservation purposes and for the scope of this analysis.

Criteria: Gross Forest Cover Loss

Gross Forest Cover Loss is defined as the area of forest cover removed because of any disturbance, including both natural and human-induced causes. Gross forest cover loss is not the same as deforestation, but a loss legal or non-regulated): swidden and permanent agriculture, artisanal logging, industrial logging, carbonization and/or fuelwood, mining, and bushfires. And indirect causes influenced by population density increasing, biophysical factors (forest fragmentation and degradation), and institutional aspects (policy failure, lack or difficulties to implement projects due to governance system, pandemics, wars and conflicts). Because at this time we cannot perform a forest cover loss analysis to spatially identify the cause of deforestation, while also considering the availability of data and aware that the forest cover gain value is not taken into account, we include all areas that suffered from forest cover loss during the period 2001-2019 among those with restoration priorities.

Parameter: CARBON CYCLE

Criteria: Carbon dioxide emissions

from tree cover loss

The carbon dioxide emission from tree cover loss reflects the emission to the atmosphere as a result of aboveground biomass loss. Emissions are gross estimates, meaning that information about the fate of land after clearing, and its associated carbon value, is not incorporated. All of the aboveground carbon is considered to be "committed" emissions to the atmosphere upon clearing. This value is associated with the annual forest cover loss data from 2001 through 2018 (from Hansen et al. 2013).

8

Criteria: Carbon accumulation potential for natural forest regrowth in reforestable areas

The carbon accumulation potential, expressed in Mg carbon/ha/yr, represents the rate at which forests could capture carbon from the atmosphere and store it in aboveground live biomass over the first 30 years of natural forest regrowth. Those values are applicable to natural forest regrowth only, and do not apply to other active restoration methods (management techniques such as planting seeds or seedlings, agroforestry, etc.). For this reason they are taken into account for their priority of conservation as potentially significant in carbon removals by natural forest regrowth.

Criteria: REDD+ project

Aware of the great potential for REDD+ projects (Reducing Emissions from Deforestation and forest Degradation) in the country, DRC has forged ahead with both the planning and implementation stages of REDD+ preparedness. REDD+ has resulted in the ongoing implementation of investment projects to tackle direct and indirect drivers of deforestation and forest degradation in the DRC. These investments include sectoral projects with a national footprint, and integrated projects at province level. US\$15 million of investments were planned in Ituri province (Kabengele, 2017) and specifically in an area south of Mambasa town, up to the border with Nord Kivu province.

For the purposes of this research, areas that are not part of already active or planned REDD+ projects are considered as priorities to encourage implementation of conservation projects in degraded or threatened areas not otherwise targeted.

Parameter: PEOPLE

Criteria: Indigenous

The protection of tropical forests is essential in order to achieve the objectives set out in the Paris Agreements (2015). This protection cannot be achieved effectively and efficiently without ensuring the participation of those most affected: the people who live in and depend on these forests. The protection of forests requires a strong local anchorage, which implies the active participation of the forest communities in the definition and implementation of development policies.

60% of the population depends directly on natural resources and food agriculture for their survival, including millions of people living in or around forests; between 400,000 and 600,000 according to established estimates (2007) are indigenous Pygmies. As such, they are particularly vulnerable to the challenges of climate change and deforestation.

The Efé are semi nomadic people who hunt and gather forest resources in the Ituri province forest for their own consumption and for exchange with sedentary subsistence farmers. Efé represent the largest group in the region, and the lands they ancestrally occupy are considered a priority for conservation action to preserve their role as land stewards.

Criteria: Displacement and Urban dynamics

Since December 2017, violence in the Ituri province has left nearly 1,000 people dead and half a million

displaced. Breaking out in the territory of Djugu, smallscale attacks first pitted the two main communities in Ituri, the Hema and Lendu, against each other. Subsequently, Lendu militias targeted the Hema, and then the national army, before attacking nearby territories. External actors, including from North Kivu province and bordering countries, are also involved.

In December 2019, the UN Office for the Coordination of Humanitarian Affairs (OCHA) estimated that 360,000 individuals had been displaced due to the latest crisis in Ituri. It acknowledged that the movement of displaced persons was continuing, and that only a portion of them (227,000) were housed in the 72 sites maintained by the UN High Commissioner for Refugees (UNHCR). Displaced people live in overcrowded makeshift camps, in dreadful conditions, and many have to travel in search of food. Continued militia attacks prevent most aid workers from reaching them and helping them protect themselves against COVID-19 pandemics and ebola outbreaks.

In addition, it should be noted that the lack of legal provisions to protect collective customary land rights encourages land seizure and eviction of groups of people without compensation and displaced in the camps.

These conflicts are creating new urban centralities that are the result of these forced migratory movements. The speed of transformation of these urban bodies is accentuated, producing a transformation of the landscape and generating clutches with the management of land resources. For this reason it is critical to plan the sustainable development of the recent urban expansion by designing an urban buffer zone that is included in this analysis as a critical area where to perform sustainability assessments.

Parameter: DRIVERS OF CONFLICT -MINING SECTOR

Criteria: Exploitation and Exploration permit

Underground minerals belong exclusively to the state. However, any private party may be authorised by the state to engage in mining activities (from exploration to exploitation and distribution), provided that specific objectives of eligibility, priority and capacity criteria set forth in the Mining Code (adopted by the Congolese Parliament on 27 January 2018) are met.

The country's subsoil is rich in minerals which are traded internationally (mining sector accounted for more than 80% of export revenues in 2015); mining is not new to the DRC but it is expanding, and exploration and exploitation permits now cover more than 39% of the country. Mining is often carried out illegally, with companies and the authorities equally failing to respect provisions of the 2018 Mining Code, including the right of mining-affected communities to compensation, and the Forest Code.

The types of mining permits available in the DRC are research/exploration permits, exploitation permits (including small-scale mines) and tailing exploitation permits. An exploration permit may be granted to any eligible private company for a period of five years, renewable once for the same duration, with respect to all mineral substances; before starting any research activity the company will have to submit a rehabilitation and mitigation plan. Should the holder of an exploration permit demonstrate through a feasibility study the existence of an economically workable ore deposit and sufficient financial capacity for the development, construction and exploitation of a mine, the Minister of Mines may grant an exploitation permit for a duration of 25 years, renewable for successive periods of 15 years. Given the limited effectiveness and feasibility of environmental management plans for areas subject to exploration and exploitation permits, those within the limits established by the Ministry of Mines are included as areas of first interest for this study. 9

Criteria: Artisanal and small-scale mine

Artisanal small-scale mining and associated infrastructure, such as construction of access roads, energy infrastructure and work camps, and inflows of mineworkers, play a great role in forest degradation. Roads fragment the forest and enable encroachment. Pressures on land use, including demand for arable land to feed mineworkers, intensifies competition for land and for areas of high conservation value forests. Ore processing contaminates water and damages other natural resources.

The development of artisanal and small-scale mining poses problems in terms of the authorities' capacity to monitor. For example, one of the biggest sources of conflict associated with mining exploitation is the uneven and opaque distribution of revenues from exploitation, and the collection of customs fees. The situation is not as critical as a few years ago (in particular from 1999 to 2003) when the management of customs and gold, wood and coffee operations had in practice fallen under the control of armed groups; although an increase in tensions would allow armed groups to take advantage of the weakness of the state to be central to trafficking operations (as revealed by field surveys carried out by IPIS in the surrounding of Mambasa observing the interference or presence of FARDC militias in the mine sites). Three representatives of local authorities described illegal taxation and ownership of gold mines by members of the 1301st Regiment of FARDC at Mongbwalu. The same sources estimated that between 15,000 and 20,000 artisanal gold diggers worked at Mongbwalu's sites during 2020.

Considering the characteristics of these extractive sites, which yield an isolated forest penetration effect on biodiversity along with their ability to generate conflict, extraction areas with processing points and auxiliary areas and their buffer zones of up to 1 km are targeted by this analysis.

Parameter: DRIVERS OF CONFLICT - OTHERS

Criteria: Artisanal logging sector and logging concession

The artisanal logging sector is characterized by a lack of clear regulations in a scenario of corruption by political and military elite; artisanal logging is descripted as lucrative but is largely informal and illegal. At the same time, the overall national framework for land allocation and forest concession zoning in the DRC is still driving deforestation and failing to protect customary land and livelihood rights. Mambasa territory is experiencing an influx of artisanal loggers from neighbouring North Kivu province and bordering countries such as Uganda and Kenya. Compared with neighboring countries, artisanal operators in eastern DRC harvest timber in dense forests; the main implication is that smallholder farmers often clear these areas once artisanal loggers have removed the largest trees.

The Forest Code issued by ministerial decree in 2002, provides that forests are state property and designates three categories of forest: classified forests, which form part of the public state domain and whose function broadly relates to environmental protection,

with highly restricted usage rights; protected forests, where local people have the right to use forest products to meet their needs in accordance with local custom; and permanent production forests, designated for exploitation through the granting of concession contracts. Artisanal permits are intended for Congolese citizens for small-scale domestic logging but are increasingly misused by local and foreign companies for logging on an industrial scale, circumventing the 2002 moratorium on new commercial concessions and enabling operators to avoid monitoring and taxes. In addition, due to a structural gap in data collection, it is hard to understand the extent of active logging concessions; therefore, for the purposes of the analysis, only industrial logging concessions are considered as the most susceptible regarding forest restoration and priority for conservation.

Criteria: Swidden and Commercial Agriculture

Rules governing access to land for agricultural purposes vary according to region and community. Despite formal legislation, access to agricultural land is generally regulated by custom. Traditional authorities such as village chiefs usually allocate land by family descent within the village. Non-descendants may rent land for food production and for fuelwood; when such cultivation ceases, the fields have to be returned to the landowner or chief. Around urban centres, producers from outside a village may rent land to farm commercially in response to demand from urban dwellers. Land may also be purchased.

Several published studies have identified swidden agriculture as one of the greatest drivers of land degradation, but in fact it is difficult to distinguish between permanent deforestation and temporary clearance for customary land use practices that allow the forest to regenerate. Agricultural production is almost entirely informal, with 90 percent of the sector driven by informal, small-scale activities.

With no detailed analysis of the different forms of agriculture, and considering the potential of some of these areas as suitable for reducing degradation through active forms of restoration such as agroecology, all areas identified as agricultural are considered as priorities for the purposes of this analysis.

Criteria: Oil concession

The oil exploration area at Lake Albert is divided into five blocks, two of which are in the Ituri province (Blocks I and II). Block III straddles the southernmost of the Ituri province and the northern part of the province of North Kivu. Explorations began in 1987 and until now have taken place discontinuously and with different impact in different blocks. There is no recent and certain information regarding the locations where exploration has already taken place, the presence of oil reserves and potentially the amount of oil that can be extracted. Evaluating the dynamics taking place in the areas subject to oil concessions on the other side of Lake Albert, in Uganda (Hoima district), where oil exploration and extraction has produced a high impact on the environment and negative repercussions on the respect of human rights, and considering that the DRC's oil sector is not legally obliged to share its revenue and contribute to local development, all the areas that fall within the perimeter of active concessions fall between those we are analyzing.



ITURI PROVINCE DEFORESTATION ALERTS 2020

37

215 A.

13

http://watetezi.org/data/DRC/analysis.html

Open the webmap to navigate and select the information

🖁 🗸 Kibali-Ituri -

Kibali-Ituri

Kibali-Ituri

-Kibali-Ituri

Kibali-Ituri

ant Homa









Parameter	Average Weight	Criteria	Subcriteria	Average Score
		Species Range-Size Rarity	> 0.0005	2
Biodiversity and natural resources	0.3	Species Richness	5-10 10-25 0.5-0.6	1 2 1
		Biodiversity Intactness Index	0.6-0-75 0.75-1	2 3
		Waterway	Waterway buffer zone 0-250 m Waterway buffer zone 250-500 m Waterway buffer zone 500 - 1000 m	3 2 1
		Protected Area	Protected Area buffer zone 0-250 m Protected Area buffer zone 250-500 m Protected Area buffer zone 500 - 1000 m	3 2 1
		Gross Forest Cover Loss	Cover loss area	3
Carbon Cycle	0.05	Carbon dioxide emissions from tree cover loss	10-25 Mg CO2 / pixel 25-50 Mg CO2 / pixel 50-100 Mg CO2 / pixel	1 2 3
		Carbon accumulation potential for natural forest regrowth in reforestable areas	1-2 Mg carbon / ha / yr 2-5 Mg carbon / ha / yr	1 2
		REDD+	Area Not included in REDD+ programs	3
		Indigenous	Presence of Indigenous people	3
People	0.2	Displacement and Urban dynamics	Inhabited area buffer zone 0-500 m Inhabited area buffer zone 500-2000 m	2 3
Conflict - Mining sector	0.25	Artisanal logging sector and logging concession	Mining site buffer zone 0-250 m Mining site buffer zone 250-500 m Mining site buffer zone 500-1000 m	3 2 1
		Exploitation and Exploration permit	Area within exploitation and exploration permit	3
Conflict - Other	0.15	Artisanal logging sector and logging concession	Area within industrial logging concession	2
		Swidden and Commercial Agriculture	Agricultural area buffer zone 0-500 m Agricultural area buffer zone 0-2000 m	2 3
		Oil concession	Area within oil concession	2
Alerts	0.05	Deforestation Alerts	Deforestation Alert Area "Confirmed"	3
		Fire Alerts	Fire Alert Area	2

• • • •

Table 1 Weights and Scores for Used Parameters

Parameter: ALERTS

Criteria: Deforestation Alerts

The GLAD alert system devised by the University of Maryland's Global Analysis and Discovery (GLAD) lab uses satellite imagery to collect weekly data on deforestation across the tropics. Each GLAD alert indicates a 30 by 30 meter area that has experienced a disturbance in the forest canopy, which indicates that trees in that area may have been lost or removed. The primary purpose of the GLAD alert system is to alert people to potential deforestation, but GLAD alerts also pick up additional disturbances such as rotation cycles in plantations, forest degradation from fires and storms, and natural changes like landslides and windthrows.

All areas where alerts were found in 2020 classified by GLAD lab as "confirmed" are included in these analyses.

Criteria: Fire Alerts

Fire alerts and thermal anomalies detected by the Visible Infrared Imaging Radiometer Suite (VIIRS), provides improved information of fire perimeters. These events (analyzed for the time frame 2013-2020) have multiple causes and they may not all have a correlation with deforestation. For instance are worth mentioning, the cultural norms in the agricultural tradition known as slash and burn (every year at the end of the dry season, farmers set fire to the bush and use the ashes as fertilizer), and on a greater scale the massive use of fire in swidden agriculture; but agricultural burning tends to lick at the edges of the dense, moist rainforests without making much of an impact. In addition, given the absence of electricity and other energy sources, open fires are widely used for cooking. As a result, increased fire density correlated with growing human density is damaging biomass with major impacts on soil fertility.

For this reason, and given the difficulty of classifying alarms according to their cause, all areas with a high fire alert density are flagged as priority for protection and conversation.

Standardization

Initially, all the criteria have diverse and dissimilar units. Standardization converts the measurement to uniform units for further processing in Weighted Overlay Analysis (WOA). The parameters lose their dimensions and the subcriteria are brought to a common suitability scale ranging from 1 to 3 in order to eliminate their inherent vagueness (Tab. 1).

RESULTS AND INTERPRETATION

The outcome of the analysis can be visualized in maps that determine the degree of sensitivity for threatened and degraded areas where to prioritize and implement conservation or active/passive restoration projects.

MAMBASA

The areas that emerged in the analysis (Fig.3) as critically threatened or degraded are those around the banks of the Ituri and Epulu rivers (in the area bordering the south of the Okapi Reserve boundary near the location of Etembo at the confluence with the Ibina River; near the locality of Teturi, on the Makeke-Mambasa axis at the intersection between the RN44 road and the river Ituri; on the Apodo-Mambasa axis at the intersection between the RP430 road and the river Epulu) and those within the Okapi reserve along the Route Nationale 4 (particularly near the localities of Bandisende, Yakele, Eboyo, Epulu).

The following case studies are reported for recognition and characterization:

Figure 4 Isolated penetrations into the forest Coordinates 719301.9, 135390.0 (EPSG:32635 WGS 84 / UTM zone 35N)

Land degradation causes: mining, logging



This site is found near the Ituri River, within dense forest. The presence of an artisanal mining site known as Apamakele is noted, and there is evidence of recent deforestation (2020) near the site. A network of logging roads was built to reach the sites at these coordinates and the many neighboring sites, up to the river banks of the Ituri River; the same river is probably used to carry people and materials to the trading centers.





Figure 5 Village along main road Coordinate 740485.0, 124807.9 (EPSG:32635 WGS 84 / UTM zone 35N)

Land degradation causes: mining, logging



The urban area of Mayuwano, on the Makeke-Mambasa road axis, saw its growth to depend on the exploitation of the adjacent mine known as Cinquantenaire (SOMIBAF) and overtime became a trading center for minerals mined in the region. Many of those who live in the area depend on the mineral trade enough to force farmers to turn into miners, attracted by the chance of a better income.

MONGBWALU

The areas that emerged in the analysis as critically threatened or degraded are those surrounded by and within the industrial mining sites of Mongbwalu City and Sayo, up to the banks of the Ituri River and still along this waterway in the south up to the administrative border with the Mambasa territories. The high amount of area degraded due to industrialscale gold mining is astonishing, and considering that a large part of the nearly depleted mineral deposits have not been subjected to any restoration project. The economy of Mongbwalu is centered on gold Figure 6 Village in the Okapi reserve Coordinate 695244.0, 149560.0 (EPSG:32635 WGS 84 / UTM zone 35N)

Land degradation causes: logging, swidden agriculture



The village of Bandisende is located along Route Nationale 4 within the Okapi reserve. The forest within this area is dominated by Gilbertiodendron dewevrei and Musanga cecropioides and is undergoing a progressive exploitation of timber by subsistence farmers, although the presence of swidden agriculture is increasing; timber sales from illegal logging are also reported to be increasing.

mining and trade. In the last century, mining in the Mongbwalu area is estimated to have produced 46,000 kg of gold; two types of gold are present in the area that also characterize the occurrence of different kinds of land degradation: l'or sec is found in flecks and small nuggets in unconsolidated mineral deposits in riverbeds or topsoil and it is obtained through open pit mining and alluvial mining, and Amalgame gold that is extracted from rocks that are obtained through underground mining and open pit mining.





The following case study is reported for recognition and characterization:

Figure 8 Industrial mining in urban area Coordinate 837652.9, 214661.5 (EPSG:32635 WGS 84 / UTM zone 35N)

Land degradation causes: artisanal and small-scale mining, industrial-scale mining



This depleted mine site is located north of Mongbwalu town, on the road that connects the town to the locality of Sayo Kati. Directly at the northern edge of this site a new high impact industrial mining area is being developed under the formal name of *Mongbwalu Gold Project*. The original morphology of the territory is totally compromised and due to the complex network of mining sites each ecological value is considered critical.

CONCLUSION

The analysis reveals that in the Mambasa and Mongbwalu sectors there are lands degraded mostly by the presence of a network of mining and related operations, which together with (illegal) logging represent a threat to the conservation of areas of high ecological interest. Policy failure, lack or difficulties to implement projects due to governance systems, the challenge of directly involving the local population, pandemics, and conflicts are among the barriers in developing conservation and restoration projects.

REFERENCES

Bailey, R.C. and Devore, I. (1989), Research on the Efe and Lese populations of the Ituri forest, Zaire. Am. J. Phys. Anthropol., 78: 459-471. https://doi.org/10.1002/ ajpa.1330780402

Bouuaert, M.C.; Zarkeyvan, M. (2018), Okapi Wildlife Reserve: Íllegal Sourcing of Gold and Timber. International Peace Information Service IPIS

Brown, E.; Makana, J.R. (2010) Experience From a Pilot Project to Improve Forest Governance in the Artisanal Logging Sector in Northeastern Democratic Republic of Congo. Wildlife Conservation Society

Cook-Patton, S.C., Leavitt, S.M., Gibbs, D. et al. (2020), Mapping carbon accumulation potential from global natural forest regrowth. Nature 585, 545-550.

Comín, F.A. (2021), Ecological Restoration and Ecosystem Services. In: Leal, F. W.; Azul, A.M.; Brandli, L.; Lange Salvia, A.; Wall, T. (eds) Life on Land. Encyclopedia of the UN Sustainable Development Goals. Springer, Cham. https:// doi.org/10.1007/978-3-319-95981-8_106

Cuvelier, J.; Marysse, S. (2004), Les enjeux économiques du conflit en Ituri, l'Afrique des Grands Lacs, 2003-2004. Annual, p. 196

DRC (2002), Law No. 007/2002 regulating the Mining Code; Loi no. 007/2002 du 11 juillet 2002 portant Code Minier

DRC (2002), Law No. 011/2002 regarding the Forest Code; Loi no. 011/2002 du 29 août 2002 portant Code Forestier

DRC (2011), Law No. 011/022 regarding basic principles for agriculture; Loi no. 011/022 du 24 décembre 2011 portant principes fondamentaux relatifs à l'agriculture

Fahey, D. (2008), Le fleuve d'or: The production and trade of gold from Mongbwalu, DRC. L'Afrique des Grands Lacs-Annuaire 2007-2008, 357-384

Hansen, M. C., P. V. Potapov, R. Moore, M. Hancher, S. A. Turubanova, A. Tyukavina, D. Thau, S. V. Stehman, S. J. Goetz, T. R. Loveland, A. Kómmareddy, A. Egorov, L. Chini, C. O. Justice, and J. R. G. Townshend. (2013), High-Resolution Global Maps of 21st-Century Forest Cover Change. Science 342 (15 November): 850-53.

Harris, N.L., Gibbs, D.A., Baccini, A. et al. (2021), Global maps of twenty-first century forest carbon fluxes. Nat. Clim. Chang. https://doi.org/10.1038/s41558-020-00976-6

Human Rights Watch (2005), The Curse of Gold. Democratic Republic of Congo. http://www.hrw.org

Ickowitz, A.; Slayback, D.; Asanzi, P.; Nasi, R. (2015), Agriculture and deforestation in the Democratic Republic of the Congo: A synthesis of the current state of knowledge. Occasional Paper 119. Bogor, Indonesia: CIFOR.

IFAD (2019), Democratic Republic of the Congo. Country Strategic Opportunities Programme 2019-2024. EB 2019/127/R.21

International Alert, Réseau Haki na Amani (2014), Oil Exploration in Ituri: A Human Rights and Conflict Risk Assessment in Block III.

International Crisis Group (2008), Congo: four priorities for sustainable peace in Ituri. Africa Report n.140

International Crisis Group (2020), DR Congo: Ending the Cycle of Violence in Ituri. Africa Report n.292

Kabengele, V. (2017), Etat d'avancement du processus national REDD+ en RDC: De la phase de préparation à la phase de paiements basés sur les résultats. Personal communication, IITA Journalist Capacity Building Workshop, 28 August 2017. Kinshasa

Kengoum, F.; Pham, T.T.; Moeliono, M.; Dwisatrio, B.; Sonwa, D.J. (2020), The context of REDD+ in the Democratic Republic of Congo: Drivers, agents and institutions. Centre for International Forestry research (CIFOR). https://doi. org/10.17528/cifor/007793

Kipalu, P.; Konè, L.; Bouchra, S.; Vig, S.; Loyombo, W. (2016), Securing Forest Peoples' Rights and Tackling Deforestation in the Democratic Republic of Congo Deforestation drivers, local impacts and rights-based solutions. Forest Peoples Programme.

Mahamba, F.; Sengenya, C. (2018), Politics and oil: the unseen drivers of violence in Congo's Ituri Province. The New Humanitarian, 4 April 2018, accessible at https://www. thenewhumanitarian.org/analysis/2018/04/04/politics-andoil-unseen-drivers-violence-congo-s-ituri-province

McHarg, I. L. (1969), Design with nature. Garden City, N.Y.: Published for the American Museum of Natural History

Newbold, T.; Hudson, L.N.; Arnell, A.P.; Contu, S. et al. (2016), Global map of the Biodiversity Intactness Index, from Newbold et al. (2016) Science. Natural History Museum Data Portal

Nyambura, R. (2017), Agrarian Transformation(s) in Africa: What's in it for Women in Rural Africa?. Development. 58. 10.1057/s41301-016-0034-0

Opelele. O.M.; Fan, W.Y.; Yu, Y.; Kachaka, S.K. (2020), Analysis of Land Use/Land Cover Change and its Prediction in the Mambasa Sector, Democratic Republic of Congo. Applied Ecology and Environmental Research, 18(4), 5627-5644.

UN Security Council (2020), Letter dated 23 December 2020 from the Group of Experts on the Democratic Republic of the Congo addressed to the President of the Security Council. S/2020/1283

Van Puijenbroek, J.; Ansoms, A. (2011), A Legacy from the Past Hindering the Future: Land Conflicts in Ituri (DRC). In: Ansoms, A.; Marysse, S. (eds) Natural Resources and Local Livelihoods in the Great Lakes Region of Africa. Palgrave Macmillan, London. https://doi. org/10.1057/9780230304994_3

Zarin, D., Harris, N.L. et al. (2016), Can carbon emissions drop by 50% in five years? Global Change Biology, 22: 1336-1347. doi:10.1111/gcb.13153

DATA

Tree cover Fraction	Hansen, M.C., Potapov, P.V., Me Thau, D., Stehman, S.V., Goetz, Justice, C.O., and Townshend, J. Forest Cover Change: Science, v content/342/6160/850.ab	
Gross Forest Cover Loss 2001-2019	Hansen, M. C., P. V. Potapov, R. Thau, S. V. Stehman, S. J. Goetz O. Justice, and J. R. G. Townsher Forest Cover Change." Science 3	
Built-up area Grid multi-temporal (1975-1990-2000-2014)	Corbane, Christina; Florczyk, At (2018): GHS built-up grid, deriv R2018A. European Commission PID: http://data.europa.eu/89h/	
Population Grid multi-temporal (1975-1990-2000-2014)	Schiavina, Marcello; Freire, Serg multitemporal (1975-1990- 200 Centre (JRC) [Dataset] doi:10.29 data.europa.eu/89h/0c6b9751-2	
Species Range-Size Rarity	IUCN 2020. The IUCN Red List o iucnredlist.org	
Species Richness	IUCN 2020. The IUCN Red List of iucnredlist.org	
Artisanal and small-scale mine	Artisanal mining site visits in E IPIS Research. https://ipisresea	
Mining Exploration Permit	Cadastre Minier - CAMI	
Mining Exploitation Permit	Cadastre Minier - CAMI	
Protected Area	Institut Congolais pour la Conse Observatory (OFAC)	
Deforestation Alert 2020	Hansen, Matthew C., Alexander Turubanova, Bryan Zutta, Suspe "Humid Tropical Forest Disturb Letters 11, no. 3 (2016): 034008.	
Fire Alerts 2013-2020	NRT VIIRS 375 m Active Fire pro earthdata.nasa.gov/firms]. doi:	
T 1 (111 (11)	TT IID T COLOR	

Industrial logging concession

Oil concession

Field crops

REDD+ pilot project areas

Area populated by Indigenous People

Tourisme (WRI/MECNT), 2013

CNREDD), 2015

Dynamique des Groupes des Peuples Autochtones (DGPA), 2012

oore, R., Hancher, M., Turubanova, S.A., Tyukavina, A., , S.J., Loveland, T.R., Kommareddy, A., Egorov, A., Chini, L., .R.G., 2013, High-Resolution Global Maps of 21st-Century v. 342, no. 6160, p. 850-853, at http://www.sciencemag.org/

Moore, M. Hancher, S. A. Turubanova, A. Tyukavina, D. , T. R. Loveland, A. Kommareddy, A. Egorov, L. Chini, C. nd. 2013. "High-Resolution Global Maps of 21st-Century 342 (15 November): 850-53

neta; Pesaresi, Martino; Politis, Panagiotis; Syrris, Vasileios ved from Landsat, multitemporal (1975-1990-2000-2014), , Joint Research Centre (JRC) doi:10.2905/jrc-ghsl-10007 jrc-ghsl-10007

gio; MacManus, Kytt (2019): GHS population grid 00-2015), R2019A. European Commission, Joint Research 005/0C6B9751-A71F-4062-830B-43C9F432370F PID: http:// a71f-4062-830b-43c9f432370f

of Threatened Species. Version 2017-3. https://www.

of Threatened Species. Version 2017-3. https://www.

astern DRC, The International Peace Information Service arch.be/home/maps-data/open-data/

ervation de la Nature (ICCN), Central African Forest

r Krylov, Alexandra Tyukavina, Peter V. Potapov, Svetlana ense Ifo, Belinda Margono, Fred Stolle, and Rebecca Moore. oance Alerts Using Landsat Data." Environmental Research . https://doi.org/10.1088/1748-9326/11/3/034008

oduct VJ114IMGTDL_NRT. Available on-line [https:// 10.5067/FIRMS/VIIRS/VJ114IMGT_NRT.002

World Resources Institute / Ministère de l'environnement Conservation de la Nature et

Ministère des Hydrocarbures (MIN.HYDRO), 2015

Thematic Agriculture Aggregation for DR Congo - AFRICOVER, FAO, 2011

Observatoire des Forêts d'Afrique Centrale / Coordination Nationale REDD (OFAC/

Environmental Defenders (ED) is an eco-feminist Environmental and Human Rights Defenders (EHRDs) organization that provide capacity-building support to Human Rights and environmental defenders in Albertine Region.

ED was established in 2017 amidst severe threats to the environment and restrictions on the work of land and environmental defenders in Uganda, specifically in the Albertine Region, West Nile and Northern Uganda, concerned for the protection of the environment and wellbeing of indigenous and rural communities who are vulnerable to poorly conceived and executed resource development projects.

ED actions are driven by the idea of protecting the environment through the conservation of forest ecology and the use of nature-based solutions as a strategy of defense against climate change. Promoting equitable governance, effective management and positive conservation outcomes. We believe that accepted environmental and human rights principles embody the right of everyone to a secure, healthy and ecologically sound environment, and that environmental degradation leads to human rights violations such as the right to life, health and culture.

